

AD-A125 608

RECOGNITION OF SURFACES IN THREE-DIMENSIONAL DIGITAL
IMAGES(U) MARYLAND UNIV COLLEGE PARK COMPUTER VISION
LAB G M REED ET AL. AUG 82 TR-1210 AFOSR-TR-83-0070


1/1


UNCLASSIFIED

AFOSR-77-3271

F/G 12/1

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

4

TR-1210
AFOSR-77-3271

August 1982

RECOGNITION OF SURFACES
IN THREE-DIMENSIONAL DIGITAL IMAGES

George M. Reed*
Azriel Rosenfeld

Computer Vision Laboratory
Computer Science Center
University of Maryland
College Park, MD 20742

S

COMPUTER SCIENCE
TECHNICAL REPORT SERIES



DTIC
ELECTE
MAR 14 1983
S B

UNIVERSITY OF MARYLAND
COLLEGE PARK, MARYLAND

20742

Approved for public release;
distribution unlimited.

83 03 14 021

UNCLASSIFIED

DTIC FILE COPY

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)
RECEIVED 100000Z JUL 82 DTIC
THIS DOCUMENT HAS BEEN REPRODUCED AS IS
AND IS NOT TO BE USED FOR ANY PURPOSE
OTHER THAN THAT FOR WHICH IT WAS
SPECIFICALLY PREPARED.
MATTHEW J. KESPER
Chief, Technical Information Division

TR-1210
AFOSR-77-3271

August 1982

RECOGNITION OF SURFACES
IN THREE-DIMENSIONAL DIGITAL IMAGES

George M. Reed*
Azriel Rosenfeld

Computer Vision Laboratory
Computer Science Center
University of Maryland
College Park, MD 20742

DTIC
ELECTE
MAR 14 1983
B

ABSTRACT

This is a continuation of a series of papers on the digital geometry of three-dimensional images. In an earlier paper by Morgenthaler and Rosenfeld, a three-dimensional analog of the two-dimensional Jordan Curve Theorem was established. This was accomplished by defining simple surface points under the symmetric consideration of 6-connectedness and 26-connectedness and by characterizing a simple closed surface as a connected collection of "orientable" simple surface points. The necessity of the assumption of orientability, a condition of often prohibitive computational cost to establish, was the major unresolved issue of that paper. In this paper, we show the assumption not to be necessary in the case of 6-connectedness and, unexpectedly, show that the property of orientability is not symmetric with respect to the two types of connectedness.

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

The support of the U.S. Air Force Office of Scientific Research under Grant AFOSR-77-3271 is gratefully acknowledged, as is the help of Janet Salzman in preparing this paper. Some of the work reported herein was accomplished while the first author was a NRC-NASA Senior Research Associate at Goddard Space Flight Center.

*Permanent address: Dept. of Mathematics, Ohio University,
Athens, Ohio 45701

1. Introduction

The digital geometry of three-dimensional images is a topic of considerable current interest, as a result of the increasing availability of 3D discrete arrays of data such as those produced in computed tomography. An introduction to the topological properties of 3D images can be found in [1].

One of the interesting problems in 3D digital topology is that of defining surfaces. Intuitively, a surface S is a set that is everywhere "thin", in the sense that in the neighborhood of any $p \in S$, there are exactly two components of \bar{S} (the complement of S), and every neighbor of $p \in S$ is adjacent to both of these components. (This definition will be stated more precisely in the next section.) In [2] it was shown that surfaces defined in this way satisfy the 3D analog of the Jordan Curve Theorem, i.e., they separate the space into two components, an "inside" and an "outside". However, this was shown only under the assumption that the surfaces were "orientable", a local property which means that for any p , the two components of \bar{S} mentioned earlier remain distinct even when we enlarge the neighborhood of p . The question of whether non-orientable surfaces exist was left open in [2].

This paper shows that when we use 6-connectedness for S (see the next section), non-orientable surfaces do not exist. Furthermore, an example is given to show that surprisingly

such surfaces can exist locally when we use 26-connectedness. However, in a subsequent paper by the first author, it will be shown that globally (i.e., if each point of the set satisfies the small neighborhood surface restriction) there can exist no 26-connected non-orientable surfaces.



Accession For	
NTIS GRAB1	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	<input type="checkbox"/>

A

2. Connectivity and simple closed surfaces

Let Σ denote a 3D array of lattice points, which, without loss of generality, we may assume to be defined by integer valued triples of Cartesian coordinates (x,y,z) . We consider two types of neighbors of a point $p = (x_p, y_p, z_p) \in \Sigma$:

(i) the neighbors (u,v,w) such that $|x_p - u| + |y_p - v| + |z_p - w| = 1$

(ii) the neighbors (u,v,w) such that $\max\{|x_p - u|, |y_p - v| + |z_p - w|\} = 1$

We refer to the neighbors of type (i) as 6-neighbors of p (the face neighbors) and to the neighbors of type (ii) as 26-neighbors of p (the face, edge, and corner neighbors). The 6-neighbors are said to be 6-adjacent to p , and the 26-neighbors are said to be 26-adjacent to p . The statement that α is a path from point p to point q in Σ means that there exists a positive integer n such that $\alpha = \{p_0, p_1, \dots, p_n\} \subseteq \Sigma$ where $p_0 = p$, $p_n = q$ and p_i is adjacent to p_{i-1} for $1 \leq i \leq n$. The terms 6-path and 26-path are utilized depending on the type of adjacency under consideration.

Let S denote a non-empty subset of Σ which, without loss of generality, we may assume does not meet the border of Σ . The points p and q of S are said to be connected in S provided there is a path from p to q which is contained in S . Connectivity is an equivalence relation, and the classes under this relation are called components. Again, the terms 6-connectivity, 26-connectivity, 6-components, and 26-components are utilized depending on the type of path under consideration.

Similarly, we can consider the components of the complement \bar{S} of S . Exactly one of these components contains the border of Σ ; this component is called the background of S . All other components of \bar{S} , if any, are called cavities in S . As is the custom in 2D digital geometry, opposite types of connectivity are assumed for S and \bar{S} to avoid anomalous situations.

Finally, let p be a point of S . We let $N_{27}(p)$ denote the 27 points in the $(3 \times 3 \times 3)$ neighborhood of p , and we let $N_{125}(p)$ denote the 125 points in the $(5 \times 5 \times 5)$ neighborhood centered at p . [Note that in [2], $N_{27}(p)$ and $N_{125}(p)$ were defined so as to exclude p . The change is made here in order to simplify the introduction of new notation.]

Surfaces

In [2], the above structure on the 3D lattice was utilized to introduce the concept of a simple closed surface in order to establish a non-trivial 3D analog of the 2D Jordan curve theorem.

A point $p \in S$ is called a simple surface point provided:

- (i) $S \cap N_{27}(p)$ has exactly one component adjacent to p (in the S sense); denote this component A_p .
- (ii) $\bar{S} \cap N_{27}(p)$ has exactly two components, C_1 and C_2 , adjacent to p (in the \bar{S} sense).
- (iii) If $q \in S$ and q is adjacent to p (in the S sense), then q is adjacent (in the \bar{S} sense) to both C_1 and C_2 .

As observed in [2], there are at most two components of $\bar{S} \cap N_{125}(p)$ adjacent (in the \bar{S} sense) to a simple surface point

p. Thus, suppose that p is a simple surface point of S and that each element of A_p is also a simple surface point of S (i.e., p is not near an "edge"). When $\bar{S} \cap N_{125}(p)$ has two components adjacent to p, (the surface at) p is said to be orientable and A_p is called a disk. When $\bar{S} \cap N_{125}(p)$ has only one component adjacent to p, (the surface at) p is said to be non-orientable and A_p is called a cross-cap.

Theorem 0. [2] If S is a connected collection of orientable simple surface points, then S has exactly one cavity, and S is said to be a simple closed surface.

We quote from [2], "(Theorem 0) is the 3D analog of the Jordan curve theorem for connected sets of simple surface points. The definition given for a simple surface point is modeled after the standard definition in continuous space, namely that a surface point is one whose neighborhood is homeomorphic with the inside of a circle on the plane. Thus every point in a small enough neighborhood of a point must be adjacent to either side of the surface.

Similarly, the concepts of orientability and cross-caps are modeled after the corresponding concepts used in the topology of continuous space. A cross-cap is homeomorphic with a Möbius strip, and may be visualized by deforming the edge of the strip to a circle in the plane. Thus, while each point on the face of the strip appears as a surface point, there is only one side (face) in the collection of points. We use the

requirement on the 125-neighborhood of a surface point to guarantee that such phenomena do not occur (at least locally).

This raises the question of the realizability of cross-caps in the 3D-lattice. That is, are the definitions of connectedness, together with the definition of simple surface point, strong enough to imply that cross-caps do not exist? From a theoretical standpoint an affirmative answer to this question would simplify the definition of simple closed surface, and from a practical viewpoint it would lessen the computational cost of detecting simple closed surfaces. While various properties such as symmetries may be used to reduce the effort needed to answer this question, the answer ultimately rests on a case analysis of the 2^{124} different configurations in the 125-neighborhood of a point $p \in S$.

3. Cross-caps in the 3D-lattice

The purpose of this paper is to answer the questions raised in [2] by (1) presenting in Example 1 a 26-connected cross-cap, and (2) establishing in Theorem 1 that there exist no 6-connected cross-caps in the 3D-lattice. Thus, we succeed in simplifying the definition of a 6-connected simple closed surface to that of a 6-connected collection of simple surface points.

Although Example 1 shows that the property of orientability is not ensured locally by the definition of a 26-connected simple surface point, the first author has established that this is so globally. It will be shown in a subsequent paper that a 26-connected collection S of points such that each point of S is a simple surface point is, in fact, orientable at each point. Hence, the assumption of orientability may be removed from Theorem 0 regardless of the type of connectedness under consideration.

Example 1. A 26-connected cross-cap.

<u>1st plane</u>	<u>2nd plane</u>	<u>3rd plane</u>	<u>4th plane</u>	<u>5th plane</u>
0 0 0 0 0	0 0 0 1 0	0 0 0 0 0	0 0 0 1 0	0 0 0 0 0
0 0 0 1 0	0 0 1 0 1	0 0 0 1 1	0 0 1 0 1	0 0 0 1 0
0 0 1 0 0	0 1 0 0 0	0 0 <u>1</u> 0 0	0 1 0 0 0	0 0 1 0 0
0 0 0 1 0	0 0 1 0 1	0 0 0 1 1	0 0 1 0 1	0 0 0 1 0
0 0 0 0 0	0 0 0 1 0	0 0 0 0 0	0 0 0 1 0	0 0 0 0 0

It should be mentioned that the deceptively simple construction of Example 1 took longer to produce than the complex proof of Theorem 1 given below. In fact, Example 1 was derived step by

step in a futile attempt to obtain a symmetric argument for 26-connectedness to that given in Theorem 1 for 6-connectedness. Perhaps the difficulty in deciding the cross-cap question was due to the fact that the answers are so counter to the intuition established by previous results. Firstly, one would expect the same answer for the two types of connectivity. Secondly, failing symmetry, one would expect that 6-connectedness, with the greater connectivity of the complement, would produce the counterexample.

Unfortunately, it is, of course, more difficult to show that something does not happen in digital geometry than to present a simple array of "0's" and "1's". The combinatorial detail involved in the following proof is unavoidable.

Theorem 1. There does not exist a 6-connected cross-cap.

3.1 Outline of the proof

Throughout the proof, p denotes a fixed simple surface point of the 6-connected subset S of Σ such that each point of $A_p = N_{27}(p) \cap S$ is also a simple surface point of S . Although it is not actually necessary to consider individually 2^{124} different configurations of $N_{125}(p)$ as suggested above, the reader of the following proof may very well begin to suspect that this is the case. Indeed, the fact that the realizability of cross-caps in the 3D-lattice depends on the choice of 6-connectedness or 26-connectedness witnesses the necessity for a detailed analysis beyond the symmetric definitions of simple surface points. The difficulty in deciding the issue for 6-connectedness ultimately focuses on (i) finding an overall strategy to upgrade the separation properties of $N_{27}(p)$ to include $N_{125}(p)$, and (ii) finding an efficient notation to describe the process.

3.2 Strategy

If M is a subset of Σ , let \bar{M} denote $M \cap \bar{S}$. Furthermore, if N is a subset of Σ which contains p , let $P(N)$ denote the property that \bar{N} has two 26-components which are 26-adjacent to p . Hence, our goal is to establish inductively that $P(N)$ holds where $N = N_{125}(p)$.

0. If $N = N_{27}(p)$, $P(N)$ holds.
1. Lemma 1 establishes that if N is the union of the top two (3×3) -planes of $N_{27}(p)$ or the union of the bottom two (3×3) -planes of $N_{27}(p)$, then \bar{N} can not have three components adjacent to p .
2. Lemma 2 uses Lemma 1 to establish that $P(N)$ holds where N is the union of five (3×3) -planes centered on p .
3. Lemma 3 uses geometric symmetry on Lemma 2 to observe that $P(N)$ holds where N is the union of three (5×3) -planes centered on p .
4. Lemma 4 uses Lemma 3 to establish that $P(N)$ holds where N is the union of five (5×3) -planes centered on p .
5. Lemma 5 uses geometric symmetry on Lemma 4 to observe that $P(N)$ holds where N is the union of three (5×5) planes centered on p .
6. Lemma 6 uses Lemma 5 to establish that $P(N)$ holds where $N = N_{125}(p)$ = the union of five (5×5) -planes centered on p .

Lemmas 2, 4, and 6 are each proved in a similar manner, but with escalating complexity, by assuming the negation and arriving at a contradiction to Lemma 1. Lemmas 2', 4', and 5' are necessary technicalities.

3.3 Notation

For each $a = (x_a, y_a, z_a)$, let $a(i, j, k) = (x_a + i, y_a + j, z_a + k)$.

In addition, let a^+ denote $a(0, 0, 1)$ and a^- denote $a(0, 0, -1)$.

For ℓ, m, n odd positive integers and k an integer, let:

$$(1) N_{m,n}^k(a) = \{a(i, j, k) \mid -\frac{(m-1)}{2} \leq i \leq \frac{(m-1)}{2} \text{ and } -\frac{(n-1)}{2} \leq j \leq \frac{(n-1)}{2}\},$$

$$(2) N_{m,n}^{k=g,h}(a) = \bigcup_{k=g,h} N_{m,n}^k(a),$$

$$(3) N_{m,n,\ell}^*(a) = N_{m,n}^{k=-\frac{(\ell-1)}{2}, \frac{(\ell-1)}{2}}(a)$$

$$(4) N_a = N_{3,3,3}^*(a), \quad N_a^k = N_{3,3}^k(a), \\ N_a^{k=g,h} = N_{3,3}^{k=g,h}(a), \text{ etc.}$$

For example:

$$N_a = N_{3,3,3}^*(a) = N_{27}(a)$$

$$N_{5,5,5}^*(a) = N_{125}(a)$$

$$N_{5,3}^{-2}(a) = \text{the } (5 \times 3)\text{-plane centered on } a(0, 0, -2)$$

$$\bar{N}_{5,3}^{-2}(a) = N_{5,3}^{-2}(a) \cap \bar{S}$$

$$N_{5,5,3}^*(a) = \text{three } (5 \times 5)\text{-planes centered on } a$$

Finally,

"#" denotes "contradiction"

"W.L.G." denotes "without loss of generality"

3.4 Proof

Lemma 1. If a is a simple surface point of S , then two 26-components of \bar{N}_a^0 cannot be merged in \bar{N}_a . Furthermore, neither of $\bar{N}_a^{i=0,1}$ nor $\bar{N}_a^{i=-1,0}$ has three 26-components.

Proof: Suppose two components, C_1 and C_2 , of \bar{N}_a^0 are merged in \bar{N}_a . Let C_1' and C_2' denote the two components of \bar{N}_a , where W.L.G. $(C_1 \cup C_2) \subseteq C_1'$.

(1) [$C_2' \cap \bar{N}_{3,3}^0(a) \neq \emptyset$, and hence \bar{N}_a^0 has three components.] Suppose $C_2' \cap \bar{N}_a^0 = \emptyset$. Then both of a^+ and a^- must be in S . To see that this is true, W.L.G. suppose $a^+ \in \bar{S}$. Then a^- must be in S or else \bar{N}_a would have only one component. Hence, since a^+ is 6-adjacent to a and $C_2' \cap \bar{N}_a^0 = \emptyset$, then $C_2' \cap \bar{N}_a^{-1} \neq \emptyset$. Now, consider the two cases where (i) some element of $C_2' \cap \bar{N}_a^{-1}$ is 6-adjacent to a^- or (ii) no element of $C_2' \cap \bar{N}_a^{-1}$ is 6-adjacent to a^- . If (i), then W.L.G. let $y = a(0, -1, -1) \in C_2'$. Then y cannot be 26-adjacent to an element of \bar{S} which is 26-adjacent to a^+ . Thus, $\{a(-1, 0, 0), a(1, 0, 0), a(-1, -1, 0), a(0, -1, 0), a(1, -1, 0)\} \subseteq S$. Furthermore, since now $(C_1 \cup C_2) \cap \{a(-1, 1, 0), a(0, 1, 0), a(1, 1, 0)\} \neq \emptyset$, it follows that $a(0, 1, 0) \in S$ (or else C_1 and C_2 would be 26-adjacent in \bar{N}_a^0) and that both $a(-1, 0, -1)$ and $a(1, 0, -1)$ are in S (or else y would be 26-adjacent to $(C_1 \cup C_2) \subseteq C_1'$). However, $a(0, 1, 0)$ is 6-adjacent to a but it cannot now be 26-connected to $y \in C_2'$ in \bar{N}_a . (# to (i)) If (ii), then W.L.G. let $y = a(1, -1, -1) \in C_2' \cap \bar{N}_a^{-1}$. Then $\{a(0, -1, 0), a(1, -1, 0), a(0, -1, -1), a(1, -1, -1)\} \subseteq S$. But, since C_1 and C_2 are not 26-connected in \bar{N}_a^0 , one of $a(-1, 0, 0)$ and $a(0, 1, 0)$ must also be in S and 6-adjacent to a . Again, this is impossible for there would not be a 6-path in \bar{N}_a from such a point to $y \in C_2'$. (# to (ii)) Thus $\{a^+, a^-\} \subseteq S$. However, since both of a^+ and a^- are 6-adjacent to a , both must be 26-adjacent to C_2' . Hence, $C_2' \cap \bar{N}_{3,3}^0(a) \neq \emptyset$. (# from which (1) follows.) Note $C_1, C_2, C_2' \cap \bar{N}_a^0$ produce three components of \bar{N}_a^0 .

(2) [The contradiction.] Since \bar{N}_a^0 has three components, three of $\{a(-1,0,0), a(0,1,0), a(0,-1,0), a(1,0,0)\}$ must be in S . Hence, W.L.G., let $\{a(-1,0,0), a(0,-1,0), a(0,1,0)\} \subseteq S$, with $y_1 = a(-1,-1,0)$ and $y_2 = a(1,-1,0)$ in different members of $\{C_1, C_2, C'_2\}$. Note that one of y_1 and y_2 is not in $(C_1 \cup C_2) \subseteq C'_1$, otherwise there could not be a 26-path in \bar{N}_a connecting $a(0,-1,0)$ to C'_2 without merging C'_1 and C'_2 . Thus, W.L.G. let $y_1 \in C_1$ and $y_2 \in C'_2$. Now, either (i) $a(0,1,0) \in C_2$, (ii) $a(0,1,0) \in S$ and $a(-1,1,0) \in C_2$, or (iii) $a(0,1,0) \in S$ and $a(1,1,0) \in C_2$. In either case, $a(-1,0,0)$ cannot be 26-adjacent to C'_2 in N_a without merging C'_1 and C'_2 . Thus, two 26-components of N_a^0 cannot be merged in \bar{N}_a . Since \bar{N}_a has two components, it follows immediately that neither of $\bar{N}_a^{i=0,1}$ or $\bar{N}_a^{i=-1,0}$ can have three components. The proof is complete.

Lemma 2. $N_{3,3,5}^*(p)$ has two components each of which is 26-adjacent to p .

Proof: (We do more work than necessary here to establish the format for the proofs of Lemmas 4 and 6.) (1) [\bar{M} has two components which are 26-adjacent to p where $M = N_p \cup N_p^2$.] Suppose not. Since \bar{N}_p has two components, C_1 and C_2 , which are 26-adjacent to p , there must exist a 26-path α in \bar{N}_p^2 from $y_1 \in N_p^1 \cap C_1$ to $y_2 \in N_p^1 \cap C_2$.

(i) [$p \in S$] Otherwise, \bar{N}_p^1 could have only one component.

(ii) [\bar{N}_{p+} has two components, C'_1 and C'_2 , which are 26-adjacent to p .] Since $p+ \in A_p$, $p+$ is also a simple surface point of S . Hence, \bar{N}_{p+} has two components 26-adjacent

to p^+ . But p is 6-adjacent to p^+ in N_{p^+} ; thus both these components must be 26-adjacent to p .

(iii) [Each of y_1 and y_2 is 26-connected to p in $\bar{N}_p^{i=0,1}$.]

Obvious.

(iv) [The contradiction from which (1) follows.] Since y_1 and y_2 are 26-connected by $\alpha \subseteq \bar{N}_p^2$, $\{y_1, y_2\}$ is contained in C_1' or C_2' , say C_1' . However, now $C_1 \cap N_p^{i=0,1}$, $C_2 \cap N_p^{i=0,1}$, and $C_2' \cap N_p^{i=0,1}$ produce three components of $\bar{N}_p^{i=0,1}$ in contradiction to Lemma 1.

(2) [$\bar{N}_{3,3,5}^* = \bar{MUN}_p^{-2}$ has two components which are 26-adjacent to p .]

This statement follows immediately from a symmetric argument to that given in (1).

Lemma 2'. Suppose a is a simple surface point of S and each of a^+ and a^- is either in \bar{S} or is also a simple surface point of S ; then $\bar{N}_{3,3,5}^*(a)$ has two components which are 26-adjacent to a .

Proof: This follows immediately from the proof of Lemma 2.

Lemma 3. $\bar{N}_{5,3,3}^*(p)$ has two components which are 26-adjacent to p .

Proof: Geometric symmetry to Lemma 2.

Lemma 3'. Consider N_a , where a is a simple surface point of S , and $b=a(-1,0,0)$, $b_1=a(-1,0,1)$, $b_2=a(0,0,-1)$ are all in S , and $D \cap C_1 \neq \emptyset$ where C_1 and C_2 are the two components of \bar{N}_a and $D =$

$\{a(1,1,0), a(1,0,0), a(1,-1,0)\}$. Then each of b and b_1 is 26-adjacent to $C_1 \cap M$, where $M = N_a^{i=-1,0}$.

Proof: Suppose N_a and M are as above and b is not 26-adjacent to $C_1 \cap M$. (Observe that b_1 is 26-adjacent to $C_1 \cap M$ if and only if b is 26-adjacent to $C_1 \cap M$.) Let $H = \{a(-1,1,0), a(0,1,0), a(1,1,0), a(-1,1,-1), a(0,1,-1), a(1,1,-1)\}$ and $K = \{a(-1,-1,0), a(0,-1,0), a(1,-1,0), a(-1,-1,-1), a(0,-1,-1), a(1,-1,-1)\}$. Since $\{a, b, b_1, b_2\} \subseteq S$ and $D \cap C_1 \neq \emptyset$, observe that if $H \cap C_2 \neq \emptyset$ and $K \cap C_2 \neq \emptyset$, there is no 26-path in $M \cap C_2$ from $H \cap C_2$ to $K \cap C_2$.

- (1) [Claim: $H \cap C_2 \neq \emptyset$ and $K \cap C_2 \neq \emptyset$] Suppose $H \cap C_2 = \emptyset$. By assumption b is not 26-adjacent to $C_1 \cap M$. But since b is 6-adjacent to a , there must exist a path α in $N_a^1 \cap C_1$ from b to $D \cap C_1$. Also, since $H \cap C_2 \neq \emptyset$, $x = a(0,1,0) \in S$. Hence, x must be 26-adjacent to C_2 . But since $D \cap C_1 \neq \emptyset$, $C_2 \cap \{a(1,0,0), a(1,0,-1), a(1,0,1)\} = \emptyset$ and it follows that x must be 26-adjacent to some $x_1 \in N_a^1 \cap C_2$. Now, since $C_1 \cap N_a^1 \neq \emptyset$ and $C_2 \cap N_a^1 \neq \emptyset$, $a(0,0,1) \in S$. Therefore, $a(0,-1,1) \in \alpha$; this is the only remaining possibility for a 26-path from b to $D \cap C_1$. Hence, $x_2 = a(0,-1,0) \notin C_2$ since it is 26-adjacent to a , and $x_2 \notin C_1$ since b is not 26-adjacent to $C_1 \cap M$. Thus, $x_2 \in S$, and being 6-adjacent to a , x_2 must be 26-adjacent to C_2 . Since $a(0,-1,1) \in C_1$, x_2 must be 26-adjacent to some point x_3 in $N_a^{-1} \cap K \cap C_2$. However, now there is no possible path in $N_a \cap C_2$ from x_1 to x_3 .# Hence $H \cap C_1 \neq \emptyset$, and similarly $K \cap C_2 \neq \emptyset$.

(2) [The contradiction.] From the above, it follows that $M \cap C_1$, $H \cap C_2$, and $K \cap C_2$ produce at least three components of $\bar{N}_a^{i=-1,0}$ in contradiction to Lemma 1.

Lemma 4. $\bar{N}_{5,3,5}^*(p)$ has two components which are 26-adjacent to p .

Proof: (1) [\bar{M} has two components which are 26-adjacent to p where $M = N_{5,3,3}^*(p) \cup N_{5,3}^2(p)$.] Suppose not. From Lemma 3, $\bar{N}_{5,3,3}^*(p)$ has 2 components, C_1 and C_2 , which are 26-adjacent to p . Thus if \bar{M} has only one component 26-adjacent to p , there must exist a 26-path α in $\bar{N}_{5,3}^2(p)$ from $y_1 \in C_1 \cap N_{5,3}^1(p)$ to $y_2 \in C_2 \cap N_{5,3}^1(p)$.

(i) [$p+ \in S$.] Suppose $p+ \in \bar{S}$. Let $a_1 = p(-1,0,1)$ and $a_2 = p(1,0,1)$.

Now $p+$ must be in one of C_1 and C_2 , say $p+ \in C_1$. Then either (a) $\{y_1, y_2\}$ is contained in the leftmost column of $N_{5,3}^1(p)$, (b) $\{y_1, y_2\}$ is contained in the rightmost column of $N_{5,3}^1(p)$, or (c) α is 26-adjacent to $p+$ and we can assume $p+ = y_1$. In either case, it follows that $\{y_1, y_2\} \subseteq \bar{N}_a^0$ and y_1 is 26-connected to y_2 in \bar{N}_a^1 where $a = a_1$ or $a = a_2$. However, since y_1 and y_2 are not 26-connected in \bar{M} , $a \in N_p \cap S$, and hence a is a simple surface point of S . But $\bar{N}_a^{i=-1,0} \subseteq \bar{M}$, and the above situation violates Lemma 1. (# from which (i) follows).

(ii) [$\bar{N}_{5,3,3}^*(p+)$ has two components, C_1' and C_2' , which are 26-adjacent to p .] Since $p+$ is a simple surface point of S and each of $p+(-1,0,0)$ and $p+(1,0,0)$ is either in \bar{S} or is a simple surface point of S , it follows from

geometric symmetry to Lemma 2' that $\bar{N}_{5,3,3}^*(p+)$ has two such components which are 26-adjacent to $p+$. Furthermore, since p is 6-adjacent to $p+$ in N_{p+} , each of these two components must be 26-adjacent to p .

(iii) [Each of y_1 and y_2 is 26-connected to p in $\bar{N}_{5,3}^{i=0,1}(p)$.] Consider y_1 arbitrarily. If $y_1 \in N_{p+}^0$, we are finished. Hence, W.L.G. let y_1 be in the leftmost column of $N_{5,3}^1(p)$. If y_1 is not 26-connected to p in \bar{M} , then $p(1,0,1)$ and $p(1,0,0)$ must be in S . However, if we assign $a=p(1,0,1)$, $b=p+$, $b_1=p$, and $b_2=p(1,0,0)$ and then apply Lemma 3' with $y_1 \in D \cap C_1$, it follows that y_1 is 26-connected to p in $\bar{N}_{p(1,0,1)}^{i=-1,0} \subseteq \bar{N}_{5,3}^{i=0,1}(p)$. (# from which (iii) follows.)

(iv) [The contradiction from which (1) follows.] Since y_1 and y_2 are 26-connected in $\bar{N}_{5,3,3}^*(p+)$, W.L.G. let $\{y_1, y_2\} \subseteq C_1'$. Now, p must be 26-adjacent to $C_2' \cap \bar{N}_p^{i=0,1}$. However, $C_1' \cap \bar{N}_p^{i=0,1}$, $C_2' \cap \bar{N}_p^{i=0,1}$, and $C_2' \cap \bar{N}_p^{i=0,1}$ produce at least three components of $\bar{N}_p^{i=0,1}$ in contradiction to Lemma 1.

(2) It now follows immediately by a symmetric argument to that given above that $\bar{N}_{3,3,5}^*(p) = \bar{M} \cup \bar{N}_{5,3}^{-2}(p)$ has two components which are 26-adjacent to p .

Lemma 4'. Suppose a is a simple surface point of S and each of $\{a(-1,0,i), a(0,0,i), a(1,0,i) \mid i=0,3\}$ is either in \bar{S} or is also a simple surface point of S ; then $\bar{N}_{5,3,5}^*(a+)$ has two components which are 26-adjacent to a .

Proof: Suppose not. Note that $\bar{N}_{5,3,3}^*(a)$ has two components 26-adjacent to a by geometric symmetry to Lemma 2', and then so does $\bar{N}_{5,3}^{i=-1,2}(a) = \bar{N}_{5,3,3}^*(a) \cup \bar{N}_{5,3}^0(a(0,0,2))$ by the proof of Lemma 4. Furthermore, if $a+ \in S$ or $a(0,0,2) \in \bar{S}$, then the proof is finished as in the proof of Lemma 4. Thus, suppose $a+ \in \bar{S}$, $a(0,0,2) \in S$, and there is a 26-path α in $\bar{N}_{5,3}^1(a(0,0,2))$ from $y_1 \in C_1 \cap \bar{N}_{5,3}^0(a(0,0,2))$ to $y_2 \in C_2 \cap \bar{N}_{5,3}^0(a(0,0,2))$ where C_1 and C_2 are the two components of $\bar{N}_{5,3}^{i=-1,2}(a)$ 26-adjacent to a . W.L.G. let $a+ \in C_1$. Then y_2 must be in either the rightmost or leftmost column of $N_{5,3}^0(a(0,0,2))$, say rightmost. Furthermore, $\{a(1,0,2), a(1,0,1)\} \subseteq S$, and by Lemma 1, y_2 and $a+$ are in opposite components of $\bar{N}_{a(1,0,2)}$. However, $a(0,0,2)$ is 6-adjacent to $a(1,0,2)$ and, by geometric symmetry to Lemma 2', $\bar{N}_{5,3,3}^*(a(0,0,2))$ has two components 26-adjacent to $a(0,0,2)$. Hence, $y_1 \in \bar{N}_{a(1,0,2)}^0$ and y_2 and $a+$ are also in opposite components of $\bar{N}_{5,3,3}^*(a(0,0,2))$. Therefore y_1 and $a+$ are in opposite components of $\bar{N}_{5,3,3}^*(a(0,0,2))$, y_1 is in the leftmost column of $N_{5,3}^0(a(0,0,2))$, $\{a(-1,0,1), a(-1,0,2)\} \subseteq S$, and y_1 and $a+$ are in opposite components of both $\bar{N}_{a(-1,0,2)}$ and $\bar{N}_{a(-1,0,1)}$. Now, since there is a path in $\bar{N}_{5,5,3}^{i=-1,2}(a)$ from a to y_1 , we have y_1 26-connected to y_1' in $\bar{N}_{a(-1,0,1)}^{i=-1,0}$. Thus, $a(-1,0,0) \in S$, and y_1' and $a+$ are in opposite components of $\bar{N}_{a(-1,0,0)}$. However, it now follows by geometric symmetry to Lemma 2' that y_1' and $a+$ are in opposite components of $\bar{N}_{5,3,3}^*(a)$ and thus of $\bar{N}_{5,3}^{i=-1,2}(a)$. Since $\{a+, y_1'\} \subseteq C_1$, this is a contradiction from which the Lemma follows. Hence $\bar{N}_{5,3}^{i=-1,3}(a) = \bar{N}_{5,3,5}^*(a+)$ has two components 26-adjacent to a .

Lemma 5. $\bar{N}_{5,5,3}^*(p)$ has two components 26-adjacent to p .

Proof: Geometric symmetry to Lemma 4.

Lemma 5'. Suppose a and b are simple surface points of S , $\bar{N}_{5,5,3}^*(a)$ has two components, C_1 and C_2 , 26-adjacent to a , and $b \in N_a^0$ is 6-connected to a via simple surface points in N_a^0 ; then $C_1' \subseteq C_1$ and $C_2' \subseteq C_2$ where C_1' and C_2' are the two components of \bar{N}_b 26-adjacent to b .

Proof: Follows immediately from the definition of simple surface point.

Lemma 6. $\bar{N}_{5,5,5}^*(p)$ has two components which are 26-adjacent to p .

Proof: (1) $[\bar{M}]$ has two components which are 26-adjacent to p where $M = N_{5,5,3}^*(p) \cup N_{5,5}^2(p)$.] Suppose not. $\bar{N}_{5,5,3}^*(p)$ has two components, C_1 and C_2 , 26-adjacent to p by Lemma 5. Hence, if \bar{M} has only one component 26-adjacent to p , there must exist a path α in $\bar{N}_{5,5}^2(p)$ from $y_1 \in C_1 \cap N_{5,5}^1(p)$ to $y_2 \in C_2 \cap N_{5,5}^1(p)$.

(i) [$p \in S$.] Suppose $p \in \bar{S}$. Then one of y_1 and y_2 cannot be 26-connected to p in $\bar{N}_{5,5}^{i=0,1}(p)$ or else y_1 would be 26-connected to y_2 in $\bar{N}_{5,5,3}^*(p)$ via p . Furthermore one of y_1 and y_2 must be 26-connected to p in $\bar{N}_{5,5}^{i=0,1}(p)$. If not, there would exist $\{q_1, q_2\} \subseteq A_p \cap N_p^0$ such that each of q_1 and q_2 is 6-connected to p in $N_p^0 \cap S$, y_1 and p are in opposite components of \bar{N}_{q_1} , and

y_2 and $p+$ are in opposite components of \bar{N}_{q_2} . But since $\bar{N}_{q_1} \cup \bar{N}_{q_2} \subset \bar{N}_{5,5,3}^*(p)$, this contradicts Lemma 5'. [To see that there exist such points q_1 and q_2 , suppose that y_1 is not 26-connected to p in $\bar{N}_{5,5}^{i=0,1}(p)$. If y_1 is not contained in $Z = \{p(-2,-2,1), p(-2,2,1), p(2,-2,1), p(2,2,1)\}$ then there must exist $q_1 \in N_p^0 \cap S$ which is 6-adjacent to p such that $y_1 \in \bar{N}_{q_1}^1$, or else y_1 would be 26-connected to $p+$ in $\bar{N}_{5,5}^{i=0,1}(p)$. Also, since y_1 and $p+$ are in different components of $\bar{N}_{q_1}^{i=0,1}$ by Lemma 1, they are in opposite components of \bar{N}_{q_1} . If y_1 is contained in Z , then W.L.G. let $y_1 = p(2,2,1)$. Again, $q_1 = p(1,1,0)$ and $x = p(1,1,1)$ must be in S , or else y_1 would be 26-connected to $p+$ in $\bar{N}_{5,5}^{i=0,1}$. Now, if both of $p(0,1,0)$ and $p(1,0,0)$ were in \bar{S} , p would be in $N_x \cap S$ but not 6-connected to x in A_x . Hence, since x is a simple surface point of S , it follows that one of $p(0,1,0)$ and $p(1,0,0)$ must be in S . Thus, q_1 is 6-connected to p in $N_p^0 \cap S$, and, as before, y_1 and $p+$ are in opposite components of \bar{N}_{q_1} . The existence of q_2 is established in a similar manner, if one supposes y_2 is not 26-connected to p in $\bar{N}_{5,5}^{i=0,1}(p)$.] Hence, W.L.G. assume y_1 is not 26-connected to p in $\bar{N}_{5,5}^{i=0,1}(p)$ but that y_2 is so connected. Now, consider y_1 . As above, if (a) y_1 is 26-connected to $y_1' \in \bar{N}_{5,5}^1(p)/Z$ in $\bar{N}_{5,5}^1(p)$, then there must exist $q \in A_p \cap N_p^1$ such that

q is 6-adjacent to p^+ and y_1' and p^+ are in opposite components of \bar{N}_q . If (b) y_1 is not 26-connected to $\bar{N}_{5,5}^1(p)/Z$ in $\bar{N}_{5,5}^1(p)$, then W.L.G. let $y_1 = p(2,2,1)$ and note that $\{p(1,1,1), p(1,2,1), p(2,0,1)\} \subseteq S$. Also, y_1 and p^+ are in opposite components of $\bar{N}_{p(1,1,1)}$. Since α must connect y_1 and y_2 in $\bar{N}_{5,5}^2(p)$, there exists $y_1' \in \{p(1,2,2), p(2,0,2)\} \cap \bar{S}$. W.L.G. let $y_1' = p(2,0,2)$. Then it follows that $q = p(1,0,1) \in S$ or else y_1 and p^+ would be 26-connected in $\bar{N}_{p(1,1,1)}$. Furthermore, since q is 6-adjacent to $p(1,1,1)$, y_1' and p^+ are in opposite components of \bar{N}_q by geometric symmetry to Lemma 1. Hence, either in case (a) or (b) we have that there exists $q \in A_p \cap N_p^1$ such that q is 6-adjacent to p^+ and p^+ and y_1' are in opposite components of \bar{N}_q where y_1' is 26-connected to y_1 in $\bar{N}_{5,5}^{i=1,2}(p)$. However, by geometric symmetry to Lemma 4', it then follows that $\bar{N}_{5,5,5}^*(p^+)$ must have two components 6-adjacent to q . Thus, y_1 and p^+ (hence y_1 and y_2) are not 26-connected by a path in $\bar{N}_{5,5}^2(p)$. (# from which (i) follows.)

- (ii) [Each of y_1 and y_2 is 26-connected to p in $\bar{N}_{5,5}^{i=0,1}(p)$.] Suppose arbitrarily that y_1 is not so connected to p . From geometric symmetry to Lemma 4', there exist two components, C_1' and C_2' of $\bar{N}_{5,5,3}^*(p^+)$ 26-adjacent to p^+ . Then from Lemma 3' as applied in the proof of Lemma 4, $y_1 \notin \{p(i,j,1) \mid i \in \{-1,0,1\}, j \in \{2,2\}\} \cup \{p(i,j,1) \mid i \in \{-2,2\}, j \in \{-1,0,1\}\}$. Thus W.L.G. let $y_1 = p(2,2,1)$ and observe that $q = p(1,1,0)$ and $q^+ = p(1,1,1)$ must be in S (hence, also in A_p) or else y_1 would be 26-connected to p in

$\bar{N}_{5,5}^{i=0,1}(p)$. Also, $x_1=p(1,2,1)$ and $x_2=p(2,1,1)$ must be in S or y_1 would again be 26-connected to p in $\bar{N}_{5,5}^{i=0,1}(p)$ by application of Lemma 3'. Now consider $H=\{p(0,1,1), p(1,0,1), p(0,1,0), p(1,0,0)\}$. Let A and B denote the two components of \bar{N}_{q+} 26-adjacent to $q+$ where $y_1 \in A$. [$H \cap A = \emptyset$.] If $H \cap A \neq \emptyset$, a 26-path in \bar{N}_{q+}^1 would merge two components of $\bar{N}_{q+}^{i=-1,0}$ in contradiction to Lemma 1. [$H \cap B \neq \emptyset$.] Suppose not. Then $H \subseteq S$. If $x_3=p(1,2,0) \in \bar{S}$, then $x_3 \in A$ and one of $K=\{p(0,1,2), p(0,2,2)\}$ must be in B since x_1 is in S and is 6-adjacent to $q+$. Thus $y=p(2,1,2) \in B$ since there is a 26-path from y_1 to y_2 in $\bar{N}_{555}^2(p)$. But q must be 26-adjacent to B and $x=p(2,0,0)$ is the only remaining possibility which implies that $p(2,0,1) \in S$. However, observe that now there is no path from x to K in \bar{N}_q which is impossible since $K \cap B \neq \emptyset$. Hence $x_3 \in S$. Now, by an application of Lemma 3', it follows that y_1 is 26-connected to $p(0,1,0)$ by a path in $\bar{N}_{q+}^{i=-1,0}$, which is impossible since $(H \cup \{x_3\}) \subseteq S$. Hence, $H \cap B \neq \emptyset$. Let $t_2 \in H \cap B$. Now consider N_{q+} . One of $p(0,1,1)$ and $p(1,0,1)$ must be in S , or else \bar{N}_{q+}^1 would merge two components of $\bar{N}_{q+}^{i=-1,0}$. Hence, since $q+$ is 6-connected to $p+$ in $N_{p+}^0 \cap S$, by Lemma 5', W.L.G. $A' \subseteq C_1'$ and $B \subseteq C_2'$. Note $t_2 \in C_2'$, and since \bar{N}_q^{-1} cannot merge two components of $\bar{N}_q^{i=0,1}$, $t_2 \in C_2$.

- (a) Suppose y_2 is not 26-connected to p in $\bar{N}_{5,5}^{i=0,1}(p)$.
 Then, as above, there exists $t_1 \in N_{p+}^{i=-1,0} \cap C_1$ such
 that t_1 is in one of C'_1 and C_2 and y_2 is in the
 other. Thus, since y_1 is 26-connected to y_2 in
 $\bar{N}_{5,5,3}^*(p+)$, $\{y_1, y_2\} \subseteq C'_1$ and $\{t_1, t_2\} \subseteq C'_2$. But $\{t_1, t_2\} \subseteq$
 $N_{p+}^{i=-1,0}$, and t_1 is not 26-connected to t_2 in $\bar{N}_{p+}^{i=-1,0}$.
 Hence, by Lemma 1, t_1 is not 26-connected to t_2 in
 $N_{5,5,3}^*(p+)$. (# to (a)).
- (b) Suppose y_2 is connected to p in $\bar{N}_{5,5}^{i=0,1}(p)$. Let $t_1 \in$
 $N_p^{i=0,1} \cap C_2$ such that t_1 is 26-connected to y_2 in
 $\bar{N}_{5,5}^{i=0,1}(p)$. Since $p \in S$, there exists $t_3 \in N_p^{i=0,1} \cap C_1$.
 Consider $\{t_1, t_2, t_3\} \subseteq N_p^{i=0,1}$. Note $\{t_1, t_2\} \subseteq C_2$ and
 therefore neither is 26-adjacent to $t_3 \in C_1$ in $\bar{N}_p^{i=0,1}$.
 Also, t_2 is not 26-connected to y_1 (hence to y_2) in
 $N_{5,5,3}^*(p+)$. Thus t_2 is not 26-connected to t_1 in
 $\bar{N}_p^{i=0,1}$. Hence, there are three components of $\bar{N}_p^{i=0,1}$
 in contradiction to Lemma 1. (# from which (ii) follows.)
- (iii) [$\bar{N}_{5,5,3}^*(p+)$ has two components, C'_1 and C'_2 , 26-adjacent
 to p .] As noted above, this follows immediately from
 geometric symmetry to Lemma 4'.
- (iv) [The contradiction to from which (1) follows.] Since
 y_1 and y_2 are 26-connected in $\bar{N}_{5,5,3}^*(p+)$, W.L.G. let
 $\{y_1, y_2\} \subseteq C'_1$. Now, p must be 26-adjacent to $C'_2 \cap \bar{N}_p^{i=0,1}$.
 However, $C_1 \cap \bar{N}_p^{i=0,1}$, $C_2 \cap \bar{N}_p^{i=0,1}$, and $C'_2 \cap \bar{N}_p^{i=0,1}$ produce at
 least three components of $\bar{N}_p^{i=0,1}$ in contradiction to
 Lemma 1.

(2) It now follows immediately by a symmetric argument to that given above that $N_{5,5,5}^*(p) = \overline{MUN}_{5,5}^{-2}(p)$ has two components which are 26-adjacent to p .

References

1. A. Rosenfeld, Three-dimensional digital topology, Info. Control 50, 1981, 119-127.
2. D. G. Morgenthaler and A. Rosenfeld, Surfaces in three-dimensional digital images, Info. Control, in press.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AFOSR-TR- 83 - 0070	2. GOVT ACCESSION NO. <i>AD-A125608</i>	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) RECOGNITION OF SURFACES IN THREE-DIMENSIONAL DIGITAL IMAGES		5. TYPE OF REPORT & PERIOD COVERED Technical
		6. PERFORMING ORG. REPORT NUMBER TR-1210
7. AUTHOR(s) George M. Reed Azriel Rosenfeld		8. CONTRACT OR GRANT NUMBER(s) AFOSR-77-3271
9. PERFORMING ORGANIZATION NAME AND ADDRESS Computer Vision Laboratory Computer Science Center University of Maryland College Park, MD 20742		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS <i>61102F/ 2304/A-2</i>
11. CONTROLLING OFFICE NAME AND ADDRESS Math & Info. Sciences, AFOSR/NM Bolling AFB Washington, DC 20332		12. REPORT DATE August 1982
		13. NUMBER OF PAGES 26
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Image processing Pattern recognition Digital geometry Three-dimensional images Surfaces		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) This is a continuation of a series of papers on the digital geometry of three-dimensional images. In an earlier paper by Morgenthaler and Rosenfeld, a three-dimensional analog of the two-dimensional Jordan Curve Theorem was established. This was accomplished by defining simple surface points under the symmetric consideration of 6-connectedness and 26-connectedness and by characterizing a simple closed surface as a connected collection of		

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

"orientable" simple surface points. The necessity of the assumption of orientability, a condition of often prohibitive computational cost to establish, was the major unresolved issue of that paper. In this paper, we show the assumption not to be necessary in the case of 6-connectedness and, unexpectedly, show that the property of orientability is not symmetric with respect to the two types of connectedness.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

4 -
DT